

Effects of exercise training on patients with CKD 2017

1. Agarwal R, Pappas MK. Delayed systolic blood pressure recovery following exercise as a mechanism of masked uncontrolled hypertension in chronic kidney disease. *Nephrol Dial Transplant.* 2017 Oct 1;32(10):1710-1717. doi: 10.1093/ndt/gfw266. PubMed PMID: 27422961.
2. Aoike DT, Baria F, Kamimura MA, Ammirati A, Cuppari L. Home-based versus center-based aerobic exercise on cardiopulmonary performance, physical function, quality of life and quality of sleep of overweight patients with chronic kidney disease. *Clin Exp Nephrol.* 2017 Jun 22. doi: 10.1007/s10157-017-1429-2. [Epub ahead of print] PubMed PMID: 28643119.
3. Bishop NC, Billany R, Smith AC. Participant acceptability of exercise in kidney disease (PACE-KD): a feasibility study protocol in renal transplant recipients. *BMJ Open.* 2017 Sep 24;7(9):e017494. doi: 10.1136/bmjopen-2017-017494. PubMed PMID: 28947458; PubMed Central PMCID: PMC5623462.
4. Bronas UG, Puzantian H, Hannan M. Cognitive Impairment in Chronic Kidney Disease: Vascular Milieu and the Potential Therapeutic Role of Exercise. *Biomed Res Int.* 2017;2017:2726369. doi: 10.1155/2017/2726369. Epub 2017 Apr 19. Review. PubMed PMID: 28503567; PubMed Central PMCID: PMC5414492.
5. Brüggemann AK, Mello CL, Dal Pont T, Hizume Kunzler D, Martins DF, Bobinski F, Pereira Yamaguti W, Paulin E. Effects of Neuromuscular Electrical Stimulation During Hemodialysis on Peripheral Muscle Strength and Exercise Capacity: A Randomized Clinical Trial. *Arch Phys Med Rehabil.* 2017 May;98(5):822-831.e1. doi: 10.1016/j.apmr.2016.12.009. Epub 2017 Jan 16. PubMed PMID: 28093194.
6. da Costa Rosa CS, Nishimoto DY, Freitas Júnior IF, Ciolac EG, Monteiro HL. Factors Associated With Levels of Physical Activity in Chronic Kidney Disease Patients Undergoing Hemodialysis: The Role of Dialysis Versus Nondialysis Day. *J Phys Act Health.* 2017 Sep;14(9):726-732. doi: 10.1123/jpah.2016-0715. Epub 2017 May 17. PubMed PMID: 28513257.
7. Downey RM, Liao P, Millson EC, Quyyumi AA, Sher S, Park J. Endothelial dysfunction correlates with exaggerated exercise pressor response during whole body maximal exercise in chronic kidney disease. *Am J Physiol Renal Physiol.* 2017 May 1;312(5):F917-F924. doi: 10.1152/ajprenal.00603.2016. Epub 2017 Mar 8. PubMed PMID: 28274927; PubMed Central PMCID: PMC5451552.
8. Highton PJ, Neale J, Wilkinson TJ, Bishop NC, Smith AC. Physical activity, immune function and inflammation in kidney patients (the PINK study): a feasibility trial protocol. *BMJ Open.* 2017 May 29;7(5):e014713. doi: 10.1136/bmjopen-2016-014713. PubMed PMID: 28554920.
9. Hiraki K, Shibagaki Y, Izawa KP, Hotta C, Wakamiya A, Sakurada T, Yasuda T, Kimura K. Effects of home-based exercise on pre-dialysis chronic kidney disease patients: a randomized pilot and feasibility trial. *BMC Nephrol.* 2017 Jun 17;18(1):198. doi: 10.1186/s12882-017-0613-7. PubMed PMID: 28623895; PubMed Central PMCID: PMC5474300.
10. Howden EJ, Lawley JS, Esler M, Levine BD. Potential role of endurance training in altering renal sympathetic nerve activity in CKD? *Auton Neurosci.* 2017 May;204:74-80. doi: 10.1016/j.autneu.2016.11.002. Epub 2016 Nov 25. Review. PubMed PMID: 27908698.
11. Ikizler TA, Robinson-Cohen C, Ellis C, Headley SAE, Tuttle K, Wood RJ, Evans EE, Milch CM, Moody KA, Germain M, Limkunakul C, Bian A, Stewart TG, Himmelfarb J. Metabolic Effects of Diet and Exercise in Patients with Moderate to Severe CKD: A Randomized Clinical Trial. *J Am Soc Nephrol.* 2017 Oct 16. pii: ASN.2017010020. doi: 10.1681/ASN.2017010020. [Epub ahead of print] PubMed PMID: 29038285.

12. Kiuchi MG, Chen S, Hoye NA. The effects of different physical activities on atrial fibrillation in patients with hypertension and chronic kidney disease. *Kidney Res Clin Pract.* 2017 Sep;36(3):264-273. doi: 10.23876/j.krcp.2017.36.3.264. Epub 2017 Sep 30. PubMed PMID: 28904878; PubMed Central PMCID: PMC5592894.
13. Kiuchi MG, Chen S. The effect of the physical activity on polymorphic premature ventricular complexes in chronic kidney disease. *Kidney Res Clin Pract.* 2017 Jun;36(2):167-174. doi: 10.23876/j.krcp.2017.36.2.167. Epub 2017 Jun 30. PubMed PMID: 28680824; PubMed Central PMCID: PMC5491163.
14. Kontos P, Alibhai SM, Miller KL et al. A prospective 2-site parallel intervention trial of a research-based film to increase exercise amongst older hemodialysis patients. *BMC Nephrol.* 2017 Jan 26;18(1):37. doi: 10.1186/s12882-017-0454-4. PMID: 28122510
15. Manfredini F, Mallamaci F, D'Arrigo G, Baggetta R, Bolignano D, Torino C, Lamberti N, Bertoli S, Ciurlino D, Rocca-Rey L, Barillà A, Battaglia Y, Rapanà RM, Zuccalà A, Bonanno G, Fatuzzo P, Rapisarda F, Rastelli S, Fabrizi F, Messa P, De Paola L, Lombardi L, Cupisti A, Fuiano G, Lucisano G, Summaria C, Felisatti M, Pozzato E, Malagoni AM, Castellino P, Aucella F, Abd ElHafeez S, Provenzano PF, Tripepi G, Catizone L, Zoccali C. Exercise in Patients on Dialysis: A Multicenter, Randomized Clinical Trial. *J Am Soc Nephrol.* 2017 Apr;28(4):1259-1268. doi: 10.1681/ASN.2016030378. Epub 2016 Dec 1. PubMed PMID: 27909047; PubMed Central PMCID: PMC5373448.
16. Manfredini F, Mallamaci F, D'Arrigo G et al. Exercise in Patients on Dialysis: A Multicenter, Randomized Clinical Trial. *J Am Soc Nephrol* 2017 28: 1259–1268. doi: 10.1681/ASN.2016030378
17. Master Sankar Raj V, Patel DR, Ramachandran L. Chronic kidney disease and sports participation by children and adolescents. *Transl Pediatr.* 2017 Jul;6(3):207-214. doi: 10.21037/tp.2017.06.03. Review. PubMed PMID: 28795012; PubMed Central PMCID: PMC5532201.
18. Michishita R, Matsuda T, Kawakami S, Tanaka S, Kiyonaga A, Tanaka H, Morito N, Higaki Y. The joint impact of habitual exercise and glycemic control on the incidence of chronic kidney disease (CKD) in middle-aged and older males. *Environ Health Prev Med.* 2017 Nov 6;22(1):76. doi: 10.1186/s12199-017-0683-y. PubMed PMID: 29165174; PubMed Central PMCID: PMC5674831.
19. Miele EM, Headley SAE, Germain M, Joubert J, Herrick S, Milch C, Evans E, Cornelius A, Brewer B, Taylor B, Wood RJ. High-density lipoprotein particle pattern and overall lipid responses to a short-term moderate-intensity aerobic exercise training intervention in patients with chronic kidney disease. *Clin Kidney J.* 2017 Aug;10(4):524-531. doi: 10.1093/ckj/sfx006. Epub 2017 Mar 27. PubMed PMID: 28852492; PubMed Central PMCID: PMC5570090.
20. Morishita S, Tsubaki A, Shirai N. Physical function was related to mortality in patients with chronic kidney disease and dialysis. *Hemodial Int.* 2017 Oct;21(4):483-489. doi: 10.1111/hdi.12564. Epub 2017 Apr 18. Review. PubMed PMID: 28418625.
21. Rahimimoghadam Z, Rahemi Z, Mirbagher Ajorpaz N, Sadat Z. Effects of Pilates exercise on general health of hemodialysis patients. *J Bodyw Mov Ther.* 2017 Jan;21(1):86-92. doi: 10.1016/j.jbmt.2016.05.012. PMID: 28167196
22. Rogan A, McCarthy K, McGregor G, Hamborg T, Evans G, Hewins S, Aldridge N, Fletcher S, Krishnan N, Higgins R, Zehnder D, Ting SM. Quality of life measures predict cardiovascular health and physical performance in chronic renal failure patients. *PLoS One.* 2017 Sep 14;12(9):e0183926. doi: 10.1371/journal.pone.0183926. eCollection 2017. Erratum in: *PLoS One.* 2017 Dec 5;12 (12):e0189382. PubMed PMID: 28910330; PubMed Central PMCID: PMC5598960.

23. Roshanravan B, Gamboa J, Wilund K. Exercise and CKD: Skeletal Muscle Dysfunction and Practical Application of Exercise to Prevent and Treat Physical Impairments in CKD. *Am J Kidney Dis.* 2017 Jun;69(6):837-852. doi: 10.1053/j.ajkd.2017.01.051. Epub 2017 Apr 18. Review. PubMed PMID: 28427790; PubMed Central PMCID: PMC5441955.
24. Santana DA, Poortmans JR, Dórea EL, Machado JBA, Fernandes AL, Sá-Pinto AL, Gualano B, Roschel H. Acute exercise does not impair renal function in nondialysis chronic kidney disease patients regardless of disease stage. *Am J Physiol Renal Physiol.* 2017 Aug 1;313(2):F547-F552. doi: 10.1152/ajprenal.00131.2017. Epub 2017 May 17. PubMed PMID: 28515176.
25. Sevick MA, Woolf K, Mattoo A, Katz SD, Li H, St-Jules DE, Jagannathan R, Hu L, Pompeii ML, Gangarossa L, Li Z, Sierra A, Williams SK, Goldfarb DS. The Healthy Hearts and Kidneys (HHK) study: Design of a 2×2 RCT of technology-supported self-monitoring and social cognitive theory-based counseling to engage overweight people with diabetes and chronic kidney disease in multiple lifestyle changes. *Contemp Clin Trials.* 2017 Sep 1. pii: S1551-7144(17)30075-7. doi: 10.1016/j.cct.2017.08.020. [Epub ahead of print] PubMed PMID: 28867396.
26. Small DM, Beetham KS, Howden EJ, Briskey DR, Johnson DW, Isbel NM, Gobe GC, Coombes JS. Effects of exercise and lifestyle intervention on oxidative stress in chronic kidney disease. *Redox Rep.* 2017 May;22(3):127-136. doi: 10.1080/13510002.2016.1276314. Epub 2017 Jan 11. PubMed PMID: 28075321.
27. Tikkainen-Dolenc H, Wadén J, Forsblom C, Harjutsalo V, Thorn LM, Saraheimo M, Elonen N, Tikkainen HO, Groop PH; FinnDiane Study Group. Physical Activity Reduces Risk of Premature Mortality in Patients With Type 1 Diabetes With and Without Kidney Disease. *Diabetes Care.* 2017 Dec;40(12):1727-1732. doi: 10.2337/dc17-0615. Epub 2017 Oct 16. PubMed PMID: 29038314.
28. Tsai YC, Chen HM, Hsiao SM, Chen CS, Lin MY, Chiu YW, Hwang SJ, Kuo MC. Association of physical activity with cardiovascular and renal outcomes and quality of life in chronic kidney disease. *PLoS One.* 2017 Aug 23;12(8):e0183642. doi: 10.1371/journal.pone.0183642. eCollection 2017. PubMed PMID: 28832653; PubMed Central PMCID: PMC5568323.
29. Watson EL, Viana JL, Wimbury D, Martin N, Greening NJ, Barratt J, Smith AC. The Effect of Resistance Exercise on Inflammatory and Myogenic Markers in Patients with Chronic Kidney Disease. *Front Physiol.* 2017 Jul 28;8:541. doi: 10.3389/fphys.2017.00541. eCollection 2017. PubMed PMID: 28804461; PubMed Central PMCID: PMC5532513.
30. Yang H, Wu X, Wang M. Exercise Affects Cardiopulmonary Function in Patients with Chronic Kidney Disease: A Meta-Analysis. *Biomed Res Int.* 2017;2017:6405797. doi: 10.1155/2017/6405797. Epub 2017 Aug 28. Review. PubMed PMID: 28932743; PubMed Central PMCID: PMC5592409.
31. Zelle DM, Klaassen G, van Adrichem E, Bakker SJ, Corpeleijn E, Navis G. Physical inactivity: a risk factor and target for intervention in renal care. *Nat Rev Nephrol.* 2017 Mar;13(3):152-168. doi: 10.1038/nrneph.2016.187. Epub 2017 Jan 31. Review. PubMed PMID: 28138130.

Effects of exercise training on dialysis patients 2017

1. Abdulkassir L, Egas-Kitchener S, Whibley D, Fynmore T, Jones GD. Captivating a captive audience: a quality improvement project increasing participation in intradialytic exercise across five renal dialysis units. *Clin Kidney J.* 2017 Aug;10(4):516-523. doi: 10.1093/ckj/sfw142. Epub 2017 Mar 15. PubMed PMID: 28852491; PubMed Central PMCID: PMC5569997.
2. Abreu CC, Cardozo LFMF, Stockler-Pinto MB, Esgalhado M, Barboza JE, Frauches R, Mafra D. Does resistance exercise performed during dialysis modulate Nrf2 and NF- κ B in patients with chronic kidney disease? *Life Sci.* 2017 Nov 1;188:192-197. doi: 10.1016/j.lfs.2017.09.007. Epub 2017 Sep 6. PubMed PMID: 28887058.
3. Böhm J, Monteiro MB, Andrade FP, Veronese FV, Thomé FS. Acute effects of intradialytic aerobic exercise on solute removal, blood gases and oxidative stress in patients with chronic kidney disease. *J Bras Nefrol.* 2017 Apr-Jun;39(2):172-180. doi: 10.5935/0101-2800.20170022. Epub 2017 Apr 27. English, Portuguese. Erratum in: *J Bras Nefrol.* 2017 Aug 17;:180. PubMed PMID: 28489182.
4. Brown PDS, Rowed K, Shearer J, MacRae JM, Parker K. Impact of intradialytic exercise intensity on urea clearance in hemodialysis patients. *Appl Physiol Nutr Metab.* 2017 Sep 29:1-4. doi: 10.1139/apnm-2017-0460. [Epub ahead of print] PubMed PMID: 28961405.
5. Bućar Pajek M, Pajek J. Characterization of deficits across the spectrum of motor abilities in dialysis patients and the impact of sarcopenic overweight and obesity. *Clin Nutr.* 2017 Mar 16. pii: S0261-5614(17)30099-7. doi: 10.1016/j.clnu.2017.03.008. [Epub ahead of print] PubMed PMID: 28343799.
6. Chang JH, Koo M, Wu SW, Chen CY. Effects of a 12-week program of Tai Chi exercise on the kidney disease quality of life and physical functioning of patients with end-stage renal disease on hemodialysis. *Complement Ther Med.* 2017 Feb;30:79-83. doi: 10.1016/j.ctim.2016.12.002. Epub 2016 Dec 7. PubMed PMID: 28137531.
7. Chigira Y, Oda T, Izumi M, Yoshimura T. Effects of exercise therapy during dialysis for elderly patients undergoing maintenance dialysis. *J Phys Ther Sci.* 2017 Jan;29(1):20-23. doi: 10.1589/jpts.29.20. Epub 2017 Jan 30. PubMed PMID: 28210031; PubMed Central PMCID: PMC5300797.
8. Chung YC, Yeh ML, Liu YM. Effects of intradialytic exercise on the physical function, depression and quality of life for haemodialysis patients: a systematic review and meta-analysis of randomised controlled trials. *J Clin Nurs.* 2017 Jul;26(13-14):1801-1813. doi: 10.1111/jocn.13514. Epub 2017 Mar 20. Review. PubMed PMID: 27532211.
9. Clarkson MJ, Fraser SF, Bennett PN, McMahon LP, Brumby C, Warmington SA. Efficacy of blood flow restriction exercise during dialysis for end stage kidney disease patients: protocol of a randomised controlled trial. *BMC Nephrol.* 2017 Sep 11;18(1):294. doi: 10.1186/s12882-017-0713-4. PubMed PMID: 28893206; PubMed Central PMCID: PMC5594594.
10. Cooke AB, Ta V, Iqbal S, Gomez YH, Mavrakanas T, Barré P, Vasilevsky M, Rahme E, Daskalopoulou SS. The Impact of Intradialytic Pedaling Exercise on Arterial Stiffness: A Pilot Randomized Controlled Trial in a Hemodialysis Population. *Am J Hypertens.* 2017 Nov 6. doi: 10.1093/ajh/hpx191. [Epub ahead of print] PubMed PMID: 29126178.
11. da Costa Rosa CS, Nishimoto DY, Freitas Júnior IF, Ciolac EG, Monteiro HL. Factors Associated With Levels of Physical Activity in Chronic Kidney Disease Patients Undergoing Hemodialysis: The Role of Dialysis Versus Nondialysis Day. *Phys Act Health.* 2017 Sep;14(9):726-732. doi: 10.1123/jpah.2016-0715. Epub 2017 May 17. PubMed PMID: 28513257.
12. Doyle A, Chalmers K, Chinn DJ, McNeill F, Dall N, Grant CH. The utility of whole body vibration exercise in haemodialysis patients: a pilot study. *Clin Kidney J.* 2017 Dec;10(6):822-829. doi: 10.1093/ckj/sfx046. Epub 2017 Jun 27. PubMed PMID: 29225812; PubMed Central PMCID: PMC5716154.
13. Dungey M, Young HML, Churchward DR, Burton JO, Smith AC, Bishop NC. Regular exercise during haemodialysis promotes an anti-inflammatory leucocyte profile. *Clin Kidney J.* 2017 Dec;10(6):813-821. doi: 10.1093/ckj/sfx015. Epub 2017 Mar 27. PubMed PMID: 29225811; PubMed Central PMCID: PMC5716206.

14. Frih B, Jaafar H, Mkacher W, Ben Salah Z, Hammami M, Frih A. The Effect of Interdialytic Combined Resistance and Aerobic Exercise Training on Health Related Outcomes in Chronic Hemodialysis Patients: The Tunisian Randomized Controlled Study. *Front Physiol.* 2017 May 31;8:288. doi: 10.3389/fphys.2017.00288. eCollection 2017. PubMed PMID: 28620308; PubMed Central PMCID: PMC5449721.
15. Frih B, Mkacher W, Bouzguenda A, Jaafar H, ALkandari SA, Ben Salah Z, Sas B, Hammami M, Frih A. Effects of listening to Holy Qur'an recitation and physical training on dialysis efficacy, functional capacity, and psychosocial outcomes in elderly patients undergoing haemodialysis. *Libyan J Med.* 2017 Dec;12(1):1372032. doi: 10.1080/19932820.2017.1372032. PubMed PMID: 28891419; PubMed Central PMCID: PMC5650043.
16. Frih B, Mkacher W, Jaafar H, Frih A, Ben Salah Z, El May M, Hammami M. Specific balance training included in an endurance-resistance exercise program improves postural balance in elderly patients undergoing haemodialysis. *Disabil Rehabil.* 2017 Jan 13:1-10. doi: 10.1080/09638288.2016.1276971. [Epub ahead of print] PubMed PMID: 28084833.
17. Fuhro MI, Dorneles GP, Andrade FP, Romão PRT, Peres A, Monteiro MB. Acute exercise during hemodialysis prevents the decrease in natural killer cells in patients with chronic kidney disease: a pilot study. *Int Urol Nephrol.* 2017 Nov 13. doi: 10.1007/s11255-017-1747-z. [Epub ahead of print] PubMed PMID: 29134614.
18. Greenwood S. Do We Need Tailored Physical Interventions to Improve Physical Function and Physical Activity Levels in Patients with Chronic Kidney Disease Treated with Peritoneal Dialysis? *Perit Dial Int.* 2017 Nov-Dec;37(6):595-597. doi: 10.3747/pdi.2017.00138. PubMed PMID: 29123000.
19. Headley S, Germain M, Wood R, Joubert J, Milch C, Evans E, Cornelius A, Brewer B, Taylor B, Pescatello LS. Blood pressure response to acute and chronic exercise in chronic kidney disease. *Nephrology (Carlton).* 2017 Jan;22(1):72-78. doi: 10.1111/nep.12730. PubMed PMID: 26786187.
20. Isnard-Rouchon M, Coutard C. [Exercise as a protective cardiovascular and metabolic factor in end stage renal disease patients]. *Nephrol Ther.* 2017 Dec;13(7):544-549. doi: 10.1016/j.nephro.2017.01.027. Epub 2017 Nov 7. French. PubMed PMID: 29126841.
21. Johansen KL, Dalrymple LS, Delgado C, Chertow GM, Segal MR, Chiang J, Grimes B, Kaysen GA. Factors Associated with Frailty and Its Trajectory among Patients on Hemodialysis. *Clin J Am Soc Nephrol.* 2017 Jul 7;12(7):1100-1108. doi: 10.2215/CJN.12131116. Epub 2017 Jun 2. PubMed PMID: 28576906; PubMed Central PMCID: PMC5498360.
22. Kang SH, Do JY, Jeong HY, Lee SY, Kim JC. The Clinical Significance of Physical Activity in Maintenance Dialysis Patients. *Kidney Blood Press Res.* 2017;42(3):575-586. doi: 10.1159/000480674. Epub 2017 Sep 18. PubMed PMID: 29017151.
23. Kontos P, Alibhai SM, Miller KL, Brooks D, Colobong R, Parsons T, Jassal SV, Thomas A, Binns M, Naglie G. A prospective 2-site parallel intervention trial of a research-based film to increase exercise amongst older hemodialysis patients. *BMC Nephrol.* 2017 Jan 26;18(1):37. doi: 10.1186/s12882-017-0454-4. PubMed PMID: 28122510; PubMed Central PMCID: PMC5267380.
24. Manfredini F, Mallamaci F, D'Arrigo G, Baggetta R, Bolignano D, Torino C, Lamberti N, Bertoli S, Ciurlino D, Rocca-Rey L, Barillà A, Battaglia Y, Rapanà RM, Zuccalà A, Bonanno G, Fatuzzo P, Rapisarda F, Rastelli S, Fabrizi F, Messa P, De Paola L, Lombardi L, Cupisti A, Fuiano G, Lucisano G, Summaria C, Felisatti M, Pozzato E, Malagoni AM, Castellino P, Aucella F, Abd ElHafeez S, Provenzano PF, Tripepi G, Catizone L, Zoccali C. Exercise in Patients on Dialysis: A Multicenter, Randomized Clinical Trial. *J Am Soc Nephrol.* 2017 Apr;28(4):1259-1268. doi: 10.1681/ASN.2016030378. Epub 2016 Dec 1. PubMed PMID: 27909047; PubMed Central PMCID: PMC5373448.
25. March DS, Graham-Brown MP, Young HM, Greenwood SA, Burton JO. 'There is nothing more deceptive than an obvious fact': more evidence for the prescription of exercise during haemodialysis (intradialytic exercise) is still required. *Br J Sports Med.* 2017 Sep;51(18):1379. doi: 10.1136/bjsports-2017-097542. Epub 2017 Mar 30. PubMed PMID: 28360144.

26. Painter PL, Agarwal A, Drummond M. Physical Function and Physical Activity in Peritoneal Dialysis Patients. *Perit Dial Int.* 2017 Nov-Dec;37(6):598-604. doi: 10.3747/pdi.2016.00256. Epub 2017 Sep 28. PubMed PMID: 28970364.
27. Qiu Z, Zheng K, Zhang H, Feng J, Wang L, Zhou H. Physical Exercise and Patients with Chronic Renal Failure: A Meta-Analysis. *Biomed Res Int.* 2017;2017:7191826. doi: 10.1155/2017/7191826. Epub 2017 Feb 20. Review. PubMed PMID: 28316986; PubMed Central PMCID: PMC5337868.
28. Rhee SY, Song JK, Hong SC, Choi JW, Jeon HJ, Shin DH, Ji EH, Choi EH, Lee J, Kim A, Choi SW, Oh J. Intradialytic exercise improves physical function and reduces intradialytic hypotension and depression in hemodialysis patients. *Korean J Intern Med.* 2017 Aug 25. doi: 10.3904/kjim.2017.020. [Epub ahead of print] PubMed PMID: 28838226.
29. Seefried L, Genest F, Luksche N, Schneider M, Fazeli G, Brandl M, Bahner U, A Heidland A. Efficacy and safety of whole body vibration in maintenance hemodialysis patients - A pilot study. *J Musculoskelet Neuronal Interact.* 2017 Dec 1;17(4):268-274. PubMed PMID: 29199185.
30. Shi Y, Zheng D, Zhang L, Yu Z, Yan H, Ni Z, Qian J, Fang W. Six-minute walk test predicts all-cause mortality and technique failure in ambulatory peritoneal dialysis patients. *Nephrology (Carlton).* 2017 Feb;22(2):118-124. doi: 10.1111/nep.12726. PubMed PMID: 26773829.
31. Shimoda T, Matsuzawa R, Yoneki K, Harada M, Watanabe T, Matsumoto M, Yoshida A, Takeuchi Y, Matsunaga A. Changes in physical activity and risk of all-cause mortality in patients on maintenance hemodialysis: a retrospective cohort study. *BMC Nephrol.* 2017 May 8;18(1):154. doi: 10.1186/s12882-017-0569-7. PubMed PMID: 28482880; PubMed Central PMCID: PMC5422904.
32. Small DM, Beetham KS, Howden EJ, Briskey DR, Johnson DW, Isbel NM, Gobe GC, Coombes JS. Effects of exercise and lifestyle intervention on oxidative stress in chronic kidney disease. *Redox Rep.* 2017 May;22(3):127-136. doi: 10.1080/13510002.2016.1276314. Epub 2017 Jan 11. PubMed PMID: 28075321.
33. Sutcliffe BK, Bennett PN, Fraser SF, Mohebbi M. The deterioration in physical function of hemodialysis patients. *Hemodial Int.* 2017 May 5. doi: 10.1111/hdi.12570. [Epub ahead of print] PubMed PMID: 28474859.
34. Tao X, Chow SKY, Wong FK. The effects of a nurse-supervised home exercise programme on improving patients' perceptions of the benefits and barriers to exercise: A randomised controlled trial. *J Clin Nurs.* 2017 Sep;26(17-18):2765-2775. doi: 10.1111/jocn.13798. Epub 2017 May 23. PubMed PMID: 28278361.
35. Wang AY, Sherrington C, Toyama T, Gallagher MP, Cass A, Hirakawa Y, Li Q, Sukkar L, Snelling P, Jardine MJ. Muscle strength, mobility, quality of life and falls in patients on maintenance haemodialysis: A prospective study. *Nephrology (Carlton).* 2017 Mar;22(3):220-227. doi: 10.1111/nep.12749. PubMed PMID: 26890468.
36. Williams S, Han M, Ye X, Zhang H, Meyring-Wösten A, Bonner M, Young C, Thijssen S, Marsh D, Kotanko P. Physical Activity and Sleep Patterns in Hemodialysis Patients in a Suburban Environment. *Blood Purif.* 2017;43(1-3):235-243. doi: 10.1159/000452751. Epub 2017 Jan 24. PubMed PMID: 28114147.
37. Wilschut ED, Rotmans JI, Bos EJ, van Zoest D, Eefting D, Hamming JF, van der Boga KEA. Supervised preoperative forearm exercise to increase blood vessel diameter in patients requiring an arteriovenous access for hemodialysis: rationale and design of the PINCH trial. *J Vasc Access.* 2017 Nov 15:0. doi: 10.5301/jva.5000826. [Epub ahead of print] PubMed PMID: 29148008.
38. Wong J, Davis P, Patidar A, Zhang Y, Vilar E, Finkelman M, Farrington K. The Effect of Intra-Dialytic Exercise on Inflammation and Blood Endotoxin Levels. *Blood Purif.* 2017;44(1):51-59. doi: 10.1159/000455059. Epub 2017 Feb 28. PubMed PMID: 28241125.
39. Zhang L, Luo H, Kang G, Wang W, Hu Y. The association between physical activity and mortality among patients undergoing maintenance hemodialysis. *Int J Nurs Pract.* 2017 Feb;23(1). doi: 10.1111/ijn.12505. Epub 2016 Dec 27. PubMed PMID: 28026071.
40. Zhao C, Ma H, Yang L, Xiao Y. Long-term bicycle riding ameliorates the depression of the patients undergoing hemodialysis by affecting the levels of interleukin-6 and interleukin-18. *Neuropsychiatr Dis Treat.* 2016 Dec 28;13:91-100. doi: 10.2147/NDT.S124630. eCollection 2017. PubMed PMID: 28096677; PubMed Central PMCID: PMC5207453.